

1. The low-energy effective action in string frame is given by

$$S = \frac{1}{2\kappa_0^2} \int d^{26} X \sqrt{-G} e^{-2\Phi} \left(R - \frac{1}{12} H_{\mu\nu\lambda} H^{\mu\nu\lambda} + 4\partial_\mu \Phi \partial^\mu \Phi \right) \quad (1)$$

Show that the equations of motions for $G_{\mu\nu}$, $B_{\mu\nu}$ and Φ are equivalent to the vanishing of the beta functions

$$\begin{aligned} \beta_{\mu\nu}(G) &= \alpha' R_{\mu\nu} + 2\alpha' \nabla_\mu \nabla_\nu \Phi - \frac{\alpha'}{4} H_{\mu\lambda\kappa} H_\nu{}^{\lambda\kappa} \\ \beta_{\mu\nu}(B) &= -\frac{\alpha'}{2} \nabla^\lambda H_{\lambda\mu\nu} + \alpha' \nabla^\lambda \Phi H_{\lambda\mu\nu} \\ \beta(\Phi) &= -\frac{\alpha'}{2} \nabla^2 \Phi + \alpha' \nabla_\mu \Phi \nabla^\mu \Phi - \frac{\alpha'}{24} H_{\mu\nu\lambda} H^{\mu\nu\lambda} \end{aligned}$$

2. Consider the string frame action (1) in D spacetime dimensions. Show that, when written in terms of the Einstein frame metric

$$\tilde{G}_{\mu\nu}(X) = e^{-4\tilde{\Phi}/(D-2)} G_{\mu\nu}(X)$$

the low-energy effective action becomes

$$S = \frac{1}{2\kappa^2} \int d^D X \sqrt{-\tilde{G}} \left(\tilde{R} - \frac{1}{12} e^{-8\tilde{\Phi}/(D-2)} H_{\mu\nu\lambda} H^{\mu\nu\lambda} - \frac{4}{D-2} \partial_\mu \tilde{\Phi} \partial^\mu \tilde{\Phi} \right)$$

where $\kappa^2 = \kappa_0^2 e^{2\Phi_0}$ and $\Phi = \Phi_0 + \tilde{\Phi}$.

3a. The string frame metric produced by N infinite static strings lying in the $(X^0, X^1) \equiv (t, x)$ direction is

$$ds^2 = f(r)^{-1} (-dt^2 + dx^2) + d\vec{X} \cdot d\vec{X}$$

where $\vec{X} = (X_2, \dots, X_{25})$ labels the space transverse to the string and

$$f(r) = 1 + \frac{g_s^2 N l_s^2}{r^{22}}$$

with $r^2 = \vec{X} \cdot \vec{X}$. Consider one further infinite probe string in this background, lying parallel to the others. Write down the Nambu-Goto action describing the motion of

this string. Show that in static gauge $t = R\tau$ and $x = R\sigma$, the low-energy excitations of the string are governed by the effective action,

$$L \approx T \int dt dx \left[-f(r)^{-1} + \frac{1}{2} \left(\frac{d\vec{X}}{dt} \cdot \frac{d\vec{X}}{dt} - \frac{d\vec{X}}{dx} \cdot \frac{d\vec{X}}{dx} \right) + \dots \right]$$

Interpret this result.

3b. Now include the coupling of the probe string to background B -field, which is given by

$$B_{01} = f(r)^{-1} - 1$$

Show that the probe string, suitably oriented and lying parallel to the initial strings, feels no static force.

4. Consider an open string whose ends are constrained to lie on a Dp -brane with a background field strength F_{ab} turned on. Show that the Neumann boundary conditions for the string must be replaced by

$$\partial_\sigma X^a - 2\pi\alpha' F^{ab} \partial_\tau X_b = 0$$

5a. Show that the Born-Infeld Lagrangian can be written in the form,

$$\mathcal{L}_{BI} \equiv \sqrt{\det(1 + F)} = \exp\left(\frac{1}{4} \text{tr} \ln(1 - F^2)\right)$$

where $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the field strength, 1 means the unit matrix, and we have set $2\pi\alpha' = 1$.

5b. Show that the equations of motion arising from the Born-Infeld action are equivalent to the beta function condition for the open string,

$$\beta_\sigma(F) = \left(\frac{1}{1 - F^2}\right)^{\mu\rho} \partial_\mu F_{\rho\sigma} = 0$$

Note: To do this, it will prove very useful if you can first show the following results:

$$\partial_\mu [\text{tr} \ln(1 - F^2)] = -4 \partial_\rho F_{\mu\sigma} \left(\frac{F}{1 - F^2}\right)^{\sigma\rho}$$

which requires use of the Bianchi identity for $F_{\mu\nu}$ and

$$\partial_\mu \left(\frac{F}{1 - F^2}\right)^{\mu\nu} = \left(\frac{F}{1 - F^2}\right)^{\mu\rho} \partial_\mu F_{\rho\sigma} \left(\frac{F}{1 - F^2}\right)^{\sigma\nu} + \left(\frac{1}{1 - F^2}\right)^{\mu\rho} \partial_\mu F_{\rho\sigma} \left(\frac{1}{1 - F^2}\right)^{\sigma\nu}$$

6*. Consider open strings in D -dimensional Minkowski space with endpoints that satisfy Dirichlet boundary conditions in the directions $X^{p+1}, X^{p+2}, \dots, X^{25}$ and Neumann conditions in the remaining directions.

Write down the classical mode expansion for an open string suspended between two separated, parallel Dp -branes.

Calculate the quantum ground state energy for this string. Find the critical separation at which this energy vanishes.

Show that the 1st excited state of the string contains a “W-boson”, a massive spin 1 particle charged under the two $U(1)$ gauge fields living on the branes. Show that this state becomes massless as the brane coincide in spacetime. Why does this mean a non-Abelian $U(2)$ gauge symmetry emerges? Now consider this system when the direction X^{25} is a circle of radius R . What is the dual description of this system that is obtained by a T-duality transformation?

7* Consider a closed string on $\mathbf{R}^{1,24} \times \mathbf{S}^1$, where $X^{25} \equiv X^{25} + 2\pi R$. Show that for general values of the radius R the massless states have level $N = \tilde{N} = 1$, with winding $m = 0$ and Kaluza-Klein momentum $n = 0$. Show that, from the perspective of an observer in $\mathbf{R}^{1,24}$, these states are a graviton, an anti-symmetric tensor, two scalars and two $U(1)$ vector fields.

Show that at the special radius $R = \sqrt{\alpha'}$, extra massless states emerge with $N = \tilde{N} = 0$, and $N = 1, \tilde{N} = 0$. What is the interpretation of these states?